# WHEN TO GO TO THE **ER** VS. **URGENT CARE VS. PEDIATRICIAN**

## **URGENT CARE**





MUSCLE STRAIN OR SPRAIN





MINOR CUT OR BURN

MINOR RASH OR INFECTION

SORE THROAT, FEVER OR COLD SYMPTOMS

## **PEDIATRICIAN**





**FUSSINESS OR COLIC** 



**EAR PAIN** 



DIARRHEA OR VOMITING

SORE THROAT, FEVER OR COLD SYMPTOMS

## **EMERGENCY ROOM**





**BROKEN BONE** 

MAJOR CUT OR BURN





Children's Liquid Tylenol (acetaminophen) 160mg/5mL		Amount of Medication	to Give per Dose
Weight (lbs)	Weight (kg)		
5.5 - 6.38 lbs	2.5 - 2.9 kg	1.25 mL	
6.4 - 7.7 lbs	3.0 - 3.5 kg	1.5 mL	
7.8 - 9.2 lbs	3.6 - 4.2 kg	1.75 mL	
9.3 - 10.6 lbs	4.3 - 4.8 kg	2 mL	
10.7 - 11.7 lbs	4.9 - 5.3 kg	2.25 mL	
11.8 - 12.8 lbs	5.4 - 5.8 kg	2.5 mL	
12.9 - 13.9 lbs	5.9 - 6.3 kg	2.75 mL	
14.0 - 16.3 lbs	6.4 - 7.4 kg	3 mL	
16.4 - 18.7 lbs	7.5 - 8.5 kg	3.5 mL	
18.8 - 20.9 lbs	8.6- 9.5 kg	4 mL	
21.0 - 23.3 lbs	9.6 - 10.6 kg	4.5 mL	
23.4 - 27.9 lbs	10.7 - 12.7 kg	5 mL	
Children's Liquid Motrin (ibuprofen) 100mg/5mL		Amount of Medication to Give per Dose	
Weight (lbs)	Weight (kg)		

#### How do I know if my child has a fever?

6-6.9 kg

7.0 - 7.9 kg

8.0 - 8.9 kg

9.0 - 9.9 kg

10.0 - 10.9 kg

Fever is 100.4°F and higher for babies and children of all ages.

3 mL

3.5 mL

4 mL

4.5 mL

5 mL

#### Why does my child have a fever?

13.2 - 15.3 lbs

15.4 - 17.5 lbs

17.6 - 19.7 lbs

19.8 - 21.9 lbs

22.0 - 23.9 lbs

A fever is a natural defense against infection. In some rare cases, teething may lead to a small rise in temperature. However, if the temperature is or above 101°F, parents are advised to consider other sources that may be causing the higher temperature, beyond just teething alone.

#### When should I treat for a fever?

For babies 2 months of age and under, parents are advised to first call their child's pediatrician and not to dose with any medication as the child needs to be evaluated, and possibly taken to an Emergency Department.

For children 2 months of age and older, with a temperature of 100.4°F or higher, parents/guardians can treat with acetaminophen or ibuprofen, as directed. However, if the fever does not remit, they should call their child's pediatrician.

